Subject Index

| 5HT receptors, 18–19 | Chorda tympani (CT) nerve, 103 |
|---|---|
| 8-OH-DPAT, 18–19 | effects of combined CT and GSP |
| A | transection, 130-131 |
| | stimulus discrimination, 131 |
| A-fibers, 105, 126 | unconditioned licking responses, |
| Acetylcholine, 20 | 130–131 |
| Agouti-related protein (AgRP), 85, | effects of CT transection, 115-125, |
| 88, 89 | 133–135, 137–138 |
| Alpha-2 adrenergic receptors, 21 | conditioned taste aversion, 122-124 |
| α-gustducin, 106, 140 | oral motor taste reactivity, 118–119 |
| Amiloride, 105–106, 121, 122, 123 | signal detection, 115–117, 137 |
| effectiveness of, 137, 138, 139 | sodium appetite, 121–122 |
| Androgen receptors, 43, 48, 55, 57 | stimulus discrimination, 119–120, 126, |
| Androgens, role in food hoarding, 80 | 137, 138 |
| Anesthesia, 142-144, 145 | on taste buds, 133–135 |
| palatal, 144 | unconditioned licking responses, |
| of tongue, 144 | 117–118 |
| Appetitive behaviors, 69, see also | innervation of salivary glands by, 113, 124 |
| Ingestive behavior, appetitive | regeneration of, 134–139 |
| phase | CT and GL cross-regeneration, |
| Aromatase activity, 48 | 140–142 |
| В | CT and GSP cross-regeneration, 140 |
| | in humans, 134–135 |
| Behavior, definition of, 101 | Circadian oscillators, 22–23 |
| Benzodiazepines, 5–6, 16 | Circadian pacemaker |
| Bmal gene, 9 | activity-dependent feedback effects, 6–7 |
| Body fat levels | dark pulse effects, 7–8 |
| changes in pregnancy, 81 | neurobiology of, 8–12 |
| inverse relation to food hoarding, | nonphotic effects, 5–7, 13, 14 |
| 80–86 | novelty-induced effects, 5, 16, 17, 19–20 |
| Brief-access taste tests, 110, 111, 117, 118, | phase shifting, 14 |
| 119, 130 | carbachol-induced, 20–21 |
| after GL and CT transection, 139 | clonidine-induced, 21 |
| С | histamine-induced, 21–22 |
| | serotonergic, 18 |
| C-fos gene, 13 Carbachol, 15–16, 20 | triazolam-induced, 19 |
| Caruachor, 13–10, 20 | trazolani-niducci, 19 |
| | |

| photic effects, 3-5, 13, 14 | receptors |
|--|--|
| photic entrainment, 12-13, 15, 16 | expressed by VMH afferents, 52 |
| sleep-deprivation effects, 5 | expressed by VMH neurons, 43, 44, |
| Circadian system, see also Circadian | 45–46, 52 |
| oscillators; Circadian pacemaker | spine density regulator, 48 |
| free-running, 2, 15 | Estrus cycle, effects on food hoarding, 80 |
| behavioral studies of, 3 | , |
| period affected by continuous | F |
| illumination, 4–5 | |
| Circumvallate papillae, 103, 106, 107, | Fasting, 73–74, 88 |
| 133, 136 | in photoperiod, 72 |
| $CKI\varepsilon$ gene, 9, 10 | Fish, 145 |
| Clock genes, 9 | Flank stimulation, 46 |
| Clock-controlled genes (CCGs), 9 | Flavor, 101 |
| Clonidine, 21 | FLI, see Fos-like immunoreactivity |
| Cold exposure, 72–73, 85 | Foliate papillae, 103, 106 |
| Consummatory behaviors, 69, see also | Food-carrying behavior, 70–71 |
| Ingestive behavior, consummatory | Food hoarding, 69–92 |
| phase | central control of, 86-91 |
| Corticotropin-releasing hormone | deficit hypothesis, 71, 74 |
| (CRH), 85 | early laboratory studies, 70–71 |
| Cry genes, 9, 10 | environmental influences, 71–78 |
| | cold exposure, 72–73, 85 |
| D | food shortage, 73–74 |
| D | food type, 74–75 |
| DARPP-32, 51 | foraging effort, 75–78 |
| Dendrites, 54 | photoperiod, 72 |
| Dietary wisdom, 75 | inverse relation to adiposity, 80-86 |
| Disinhibition, 143–144 | lipostatic theory, 71 |
| Dopamine receptors, 51 | in nature, 70–71 |
| Drosophila clock genes, 9 | peripheral physiological controls of, |
| | 78–80 |
| E | gonadal steroids, 79–80 |
| TO THE STATE OF TH | metabolic fuel challenges, 78–79 |
| ENaCs, see Epithelial sodium channels | Food intake, role of VMH in, 56 |
| Energy acquisition, 70 | Food restriction, 73 |
| Energy balance, 86, 92 | Foraging, 69, 87–90 |
| Energy flux, 86, 92 | effort, 75–78 |
| Energy stores, 75, 81, 85 | Fos induction, 46 |
| Enkephalin, 41, 42, 43 | Fos-like immunoreactivity (FLI), |
| Epiglottis, 106 | 127–129 |
| Epinephrine, 78, 79 | Frustration, 71 |
| Epithelial sodium channels (ENaCs), | Fungiform papillae, 102, 107, 134 |
| 105–106, 112, 121, 123, 138 | G |
| $ER\alpha$, 48, 50–51, 56 | G |
| $ER\beta$, 48, 50–51, 56 | G-protein genes, 107 |
| Estradiol, 53, 80 | GABA, 10, 11, 15, 16, 17 |
| Estrogen | in VMH, 46, 47, 48, 49, 52 |
| cellular effects of, 50-51 | GAD, 47 |
| neonatal exposure to, 49 | GAP-43, 49 |
| neuronal activity effects of, 51-52 | Gaping, 127-129, 132, 136 |
| | |

| Geniculohypothalamic tract (GHT), 15–17 Geschmacksstreifen, 103, 106, 134 GHT, see Geniculohypothalamic tract Glossopharyngeal (GL) nerve, 103 effects of GL transection, 125–129, 131 oral motor taste reactivity, 127–129 sodium appetite, 126 stimulus discrimination, 120, 125–126, 131 on taste buds, 133, 134 | early history, 113–115 general caveats, 112–113 GL transection, see Glossopharyngeal (GL) nerve, effects of GL transection summary and functional implications, 131–133 H H-fibers, 104, 105 Hibernation, 72 |
|---|--|
| unconditioned licking responses, 125 human, 144 | Histamine, 21–22 |
| innervation of von Ebner's glands by, 113 | Hunger, 71 Hyperphagia, 74 |
| regeneration of, 134–136, 139 | I |
| GL and CT cross-regeneration, | IGL, see Intergeniculate leaflet |
| 140–142 | Immediate early-response genes |
| Glucoprivation, 79 | (IEGs), 13, 16 |
| Glutamate, 13, 16, 17, 21, 22 | Infertility, fasting-induced, 83 |
| in VMH, 46, 47, 52 | Ingestive behavior |
| Goal-lessness, 71 | appetitive phase of, 69, 87, 88, 90, 111, |
| Golgi impregnation, 53 | 127, see also Food hoarding; |
| Gonadal fat, 78 | Foraging |
| Gonadal steroids, 79–80 | consummatory phase of, 69, 87, 90-91, |
| Greater superficial petrosal (GSP) | 111, 127 |
| nerve, 103, 114 | Inosine monophosphate, 107 |
| effects of combined CT and GSP | Instinct, 71 |
| transection, 130–131 | Insulin, 79, 111, 123 |
| stimulus discrimination, 131 unconditioned licking responses, | Intake tests, 110, 111, 113–115, 141 |
| 130–131 | Intergeniculate leaflet (IGL), 11, |
| regeneration of, 134 | 15–17, 18, 19 |
| GSP and CT cross-regeneration, 140 | K |
| Gustatory system, peripheral, 101–146 | Kinase C, 51 |
| anatomy, 102–103, 104 | Killase C, 31 |
| biophysics and molecular biology, 105 | L |
| electrophysiology, 103-105 | Lateral hypothalamus (LH), 85 |
| human, 142–145 | Leptin, 82–83 |
| compared with rodents, 146 | Licking responses, 110, 117–118, 119, 125, |
| nerve cross-regeneration, 140–142 | 130–131, 145 |
| nerve regeneration consequences, | Lipectomy, 81 |
| 133–139 | Lipid deficits, 81–82, 86 |
| nerve transection consequences, | sensed by brain, 82–83 |
| 111-133 | Lipoprivation, 79 |
| CT and GSP transection, see Chorda | Lordosis response, 39–40 |
| tympani (CT) nerve, effects of | estrogen mediated, 50–51 in male rats, 49 |
| combined CT and GSP | * |
| transection | role of GABA in, 46, 47 role of glutamate in, 47 |
| CT transection, see Chorda tympani (CT) nerve, effects of CT transection | role of VMH in, 44, 47, 58 |
| nerve, effects of C1 transection | 1016 Of VIVIII III, 77, 77, 50 |
| | |

| M | Periaqueductal gray (PAG), 40, 41, |
|--|--|
| Macronutrients, 75 | 42, 54 |
| Masking, 2 | role in lordosis reflex, 44 subdivisions of, 42 |
| Maternal aggression, 55 | Peripheral gustatory system, see Gustatory |
| Maternal behavior role, 55 | |
| Maternal nurturing, 55 | system, peripheral Phase–response curves (PRCs), 2 |
| Mecamylamine, 20 | for carbachol-induced phase- |
| Melanocortin (MC), 89 | shifting, 20 |
| Melanopsin, 13 | |
| Melatonin, 23, 72, 86 | dark pulse, 7–8 nonphotic, 6, 14, 20 |
| Metabolic fuel utilization, 79 | for novelty-induced activity, 4, 5, 7 |
| mGluR4 receptors, 107, 112 | |
| Micronutrients, 75 | photic, 3–4, 6, 14, 20 for serotonergic stimulation, 18 |
| MK-801, 21 | triazolam effects, 4, 5, 7 |
| Morphine, 90, 91 | |
| • | Photoperiod, 72, 86 |
| N | Pineal gland, 72 |
| N-fibers, 104, 105 | Pituitary adenyl cyclase activating peptide |
| Naltrexone, 90, 91 | (PACAP), 11, 15 Posterior palatine field, 103, 134 |
| Nasoincisor ducts (NID), 130, 134 | |
| Neuropeptide Y (NPY), 11, 15, 16, 17, 88 | PRCs, see Phase-response curves |
| injected, 85, 87, 88, 89 | Predation, 92 |
| | Progesterone, 80 |
| Neuropeptides as controllers of appetitive ingestive | Progestin receptors, 44, 46, 48 |
| behaviors, 88–90 | Prolactin, 42, 43, 80 |
| as controllers of food intake, 87 | PRV detected in, 45 Proopiomelanocortin gene, 88 |
| Neuropil, in VMH, 48 | Pseudorabies virus (PRV), 44–46 |
| NPY, see Neuropeptide Y | Pup cannibalism, 75 |
| Nucleus of the solitary tract (NST), 103, | PVN, see Paraventricular nucleus |
| 127–128, 135, 139, 143, 145 | 1 VIV, see I araventricular racicus |
| 127 120, 133, 133, 113, 115 | Q |
| 0 | |
| 0 : 4 = 00 | Q-fibers, 105, 126 |
| Opiates, 90 | Quinine aversion, 114, 115 |
| Oral motor taste reactivity, 118–119, | _ |
| 127–129 | R |
| Orexin, 88, 89 | Raphe nuclei, mesencephalic, 11, 17-20 |
| Oxytocin receptors, 43, 44, 51 | Reproduction, and food hoarding, 80, 86, 92 |
| P | Retinohypothalamic tract (RHT), |
| PAG, see Periaqueductal gray | 12–15, 17 |
| Paraventricular nucleus (PVN), 40-41 | 12 13, 17 |
| Per genes, 9, 10, 14, 17 | S |
| expression effects | 5 |
| of 8-OH-DPAT, 19 | S-fibers, 105 |
| of glutamate, 13 | Salivary glands, innervation of, 113, |
| of light exposure, 17 | 124–125 |
| of PACAP, 15 | Salivation, 111, 133 |
| expression in peripheral tissues, 23 | SCN, see Suprachiasmatic nucleus |
| functional specialization, 22 | Security, 71 |
| Perception, 101 | Serotonin, 17–20 |
| | |

| Sexual behavior human, 57 importance of VMH for, 40, 55 rodent, 39 SF-1 gene, 55 SLN, see Superior laryngeal (SLN) nerve Sodium appetite, 121–122, 126 Spines, dendritic, 53, 54, 58 Splitting, 22 Substance P (SP), 11, 15, 42, 43, 46, 56 in WAT, 83 Superior laryngeal (SLN) nerve, 103, 114, 132 minor role in taste-guided behavior, 130 Suprachiasmatic nucleus (SCN) circadian oscillation in, 8–9 core, 10–12 GRP-positive neurons in, 10 VIP-positive neurons in, 10 SCN afferent systems, 11, 12–22 IGL, 11, 15–17 raphe, 11, 17–20 retinal, 11, 12–15 shell afferents, 20–22 | Temperature rhythm, daily, 2 Testosterone, 80 neonatal exposure to, 49 Thermal homeostasis, 72 Thiouracil, 79 Thyroxine, 79 Tim gene, 9 Tongue maps, 108–109 Triazolam, 5–6, 16, 17, 19–20 Triglyceride, 111 U U-50,488H, 90, 91 Umami, 107 V Vaginal cervix stimulation (VCS), 39, 46 Ventromedial nucleus of the hypothalamus (VMH), 39–58 afferents, 40–41, 49, 52 calorie regulation role, 55–56 cellular effects of estrogen in, 50–51 central zone, 43 dorsomedial zone, 43 |
|---|--|
| shell, 10–12 AVP-positive neurons in, 10 | spine density in, 53, 54 efferents, 42 |
| Т | human, 56–57 |
| T1Rs, 107 | intrinsic circuitry, 43–47 |
| T2Rs, 106, 107 | male sexual behavior role, 55 |
| Taste, 101 | maternal behavior role, 55 neuronal activity effects of estrogen in, |
| Taste aversion, conditioned, 122–124 | |
| | · · · · · · · · · · · · · · · · · · · |
| Taste buds, 102–103, 104 | 51–52 |
| after nerve transaction, 133-134 | 51–52 rostral zone, 43, 45 |
| after nerve transaction, 133-134 regeneration of, 134, 135, 136, 139 | 51–52 |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, | 51–52 rostral zone, 43, 45 sexual dimorphisms of, 47–50 synaptic organization, 44, 48 estrogen-induced changes, 52–55 |
| after nerve transaction, 133-134 regeneration of, 134, 135, 136, 139 | 51–52 rostral zone, 43, 45 sexual dimorphisms of, 47–50 synaptic organization, 44, 48 estrogen-induced changes, 52–55 ventrolateral zone, 43, 44 |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, 122, 125–126, 131, 137–138, 144–145 | 51–52 rostral zone, 43, 45 sexual dimorphisms of, 47–50 synaptic organization, 44, 48 estrogen-induced changes, 52–55 ventrolateral zone, 43, 44 neurons in, 45–46, 54 |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, 122, 125–126, 131, 137–138, | 51–52 rostral zone, 43, 45 sexual dimorphisms of, 47–50 synaptic organization, 44, 48 estrogen-induced changes, 52–55 ventrolateral zone, 43, 44 neurons in, 45–46, 54 role in lordosis response, 44, 47, 58 |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, 122, 125–126, 131, 137–138, 144–145 Taste dysfunction, human, 142–145 Taste function, 108–109 domains of, 109–111 | sexual dimorphisms of, 47–50 synaptic organization, 44, 48 estrogen-induced changes, 52–55 ventrolateral zone, 43, 44 neurons in, 45–46, 54 role in lordosis response, 44, 47, 58 spine density in, 53–54 |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, 122, 125–126, 131, 137–138, 144–145 Taste dysfunction, human, 142–145 Taste function, 108–109 domains of, 109–111 affective, 109, 110–111 | source for the street of the s |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, 122, 125–126, 131, 137–138, 144–145 Taste dysfunction, human, 142–145 Taste function, 108–109 domains of, 109–111 affective, 109, 110–111 physiological domain, 111 | rostral zone, 43, 45 sexual dimorphisms of, 47–50 synaptic organization, 44, 48 estrogen-induced changes, 52–55 ventrolateral zone, 43, 44 neurons in, 45–46, 54 role in lordosis response, 44, 47, 58 spine density in, 53–54 Visual system, primate, 146 VMH, see Ventromedial nucleus of the |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, 122, 125–126, 131, 137–138, 144–145 Taste dysfunction, human, 142–145 Taste function, 108–109 domains of, 109–111 affective, 109, 110–111 physiological domain, 111 sensory-discriminative, 109–110 | source for the street of the s |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, 122, 125–126, 131, 137–138, 144–145 Taste dysfunction, human, 142–145 Taste function, 108–109 domains of, 109–111 affective, 109, 110–111 physiological domain, 111 sensory-discriminative, 109–110 Taste phantoms, 143 | rostral zone, 43, 45 sexual dimorphisms of, 47–50 synaptic organization, 44, 48 estrogen-induced changes, 52–55 ventrolateral zone, 43, 44 neurons in, 45–46, 54 role in lordosis response, 44, 47, 58 spine density in, 53–54 Visual system, primate, 146 VMH, see Ventromedial nucleus of the hypothalamus |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, 122, 125–126, 131, 137–138, 144–145 Taste dysfunction, human, 142–145 Taste function, 108–109 domains of, 109–111 affective, 109, 110–111 physiological domain, 111 sensory-discriminative, 109–110 Taste phantoms, 143 Taste pore, 102 | rostral zone, 43, 45 sexual dimorphisms of, 47–50 synaptic organization, 44, 48 estrogen-induced changes, 52–55 ventrolateral zone, 43, 44 neurons in, 45–46, 54 role in lordosis response, 44, 47, 58 spine density in, 53–54 Visual system, primate, 146 VMH, see Ventromedial nucleus of the hypothalamus W |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, 122, 125–126, 131, 137–138, 144–145 Taste dysfunction, human, 142–145 Taste function, 108–109 domains of, 109–111 affective, 109, 110–111 physiological domain, 111 sensory-discriminative, 109–110 Taste phantoms, 143 Taste pore, 102 Taste receptor cells, 101, 102 | rostral zone, 43, 45 sexual dimorphisms of, 47–50 synaptic organization, 44, 48 estrogen-induced changes, 52–55 ventrolateral zone, 43, 44 neurons in, 45–46, 54 role in lordosis response, 44, 47, 58 spine density in, 53–54 Visual system, primate, 146 VMH, see Ventromedial nucleus of the hypothalamus W White adipose tissues (WAT), 78, 81–82, |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, 122, 125–126, 131, 137–138, 144–145 Taste dysfunction, human, 142–145 Taste function, 108–109 domains of, 109–111 affective, 109, 110–111 physiological domain, 111 sensory-discriminative, 109–110 Taste phantoms, 143 Taste pore, 102 Taste receptor cells, 101, 102 ion channels on, 105 | rostral zone, 43, 45 sexual dimorphisms of, 47–50 synaptic organization, 44, 48 estrogen-induced changes, 52–55 ventrolateral zone, 43, 44 neurons in, 45–46, 54 role in lordosis response, 44, 47, 58 spine density in, 53–54 Visual system, primate, 146 VMH, see Ventromedial nucleus of the hypothalamus W White adipose tissues (WAT), 78, 81–82, 83–85 |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, 122, 125–126, 131, 137–138, 144–145 Taste dysfunction, human, 142–145 Taste function, 108–109 domains of, 109–111 affective, 109, 110–111 physiological domain, 111 sensory-discriminative, 109–110 Taste phantoms, 143 Taste pore, 102 Taste receptor cells, 101, 102 ion channels on, 105 laryngeal, 102, 114 | rostral zone, 43, 45 sexual dimorphisms of, 47–50 synaptic organization, 44, 48 estrogen-induced changes, 52–55 ventrolateral zone, 43, 44 neurons in, 45–46, 54 role in lordosis response, 44, 47, 58 spine density in, 53–54 Visual system, primate, 146 VMH, see Ventromedial nucleus of the hypothalamus W White adipose tissues (WAT), 78, 81–82, 83–85 epididymal (EWAT), 78, 81 |
| after nerve transaction, 133–134 regeneration of, 134, 135, 136, 139 Taste discrimination tasks, 119–120, 122, 125–126, 131, 137–138, 144–145 Taste dysfunction, human, 142–145 Taste function, 108–109 domains of, 109–111 affective, 109, 110–111 physiological domain, 111 sensory-discriminative, 109–110 Taste phantoms, 143 Taste pore, 102 Taste receptor cells, 101, 102 ion channels on, 105 | rostral zone, 43, 45 sexual dimorphisms of, 47–50 synaptic organization, 44, 48 estrogen-induced changes, 52–55 ventrolateral zone, 43, 44 neurons in, 45–46, 54 role in lordosis response, 44, 47, 58 spine density in, 53–54 Visual system, primate, 146 VMH, see Ventromedial nucleus of the hypothalamus W White adipose tissues (WAT), 78, 81–82, 83–85 |