

An analog of Titchmarsh's theorem for the Laguerre–Bessel transform

Analog of
Titchmarsh's
theorem

Larbi Rakhimi and Radouan Daher

*Department of Mathematics, Faculty of Sciences Ain Chock,
University Hassan II Casablanca, Casablanca, Morocco*

Abstract

Purpose – Using a generalized translation operator, this study aims to obtain a generalization of Titchmarsh's theorem for the Laguerre–Bessel transform for functions satisfying the Ψ -Laguerre–Bessel–Lipschitz condition in the space $L^p_{\alpha}(\mathbb{K})$, where $\mathbb{K} = [0, +\infty[\times [0, +\infty[$.

Design/methodology/approach – The author has employed the results developed by Titchmarsh, of reference number [1].

Findings – In this paper, an analogous of Titchmarsh's theorem is established for Laguerre–Bessel transform.

Originality/value – To the best of the authors' findings, at the time of submission of this paper, the results reported are new and interesting.

Keywords Laguerre–Bessel transform, Generalized translation operator, Lipschitz class, Titchmarsh theorem

Paper type Research paper

Received 25 April 2022
Revised 27 August 2022
15 October 2022
31 December 2022
Accepted 17 January 2023

1. Introduction

The integral Fourier transform, as Fourier series, is widely used in various fields of calculus, computational mathematics, mathematical physics, etc.

Years ago, Titchmarsh established ([1], Theorem 84) that if f satisfies the Lipschitz condition $Lip(\delta; p)$ in the L^p norm ($1 < p \leq 2$) on the real line \mathbb{R} , that is

$$\left(\int_{\mathbb{R}} |f(x+h) - f(x)|^p \right)^{\frac{1}{p}} = o(h^{\delta}), \quad (0 < \delta \leq 1) \quad h \rightarrow 0.$$

Then its Fourier transform $\mathcal{F}(f)$ belongs to $L^{\beta}(\mathbb{R})$, for

$$\frac{p}{p + \delta p - 1} < \beta \leq \frac{p}{p - 1}.$$

A second result ([1], Theorem 85) characterized the set of functions in $L^2(\mathbb{R})$ satisfying the Cauchy–Lipschitz condition by means of an asymptotic estimate growth of the norm of their Fourier transform. Namely, we have:

JEL Classification — 26A16, 43A30, 26D10

© Larbi Rakhimi and Radouan Daher. Published in *the Arab Journal of Mathematical Sciences*. Published by Emerald Publishing Limited. This article is published under the Creative Commons Attribution (CC BY 4.0) license. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this license may be seen at <http://creativecommons.org/licenses/by/4.0/legalcode>

The authors would like to thank the referee for carefully checking the manuscript and for his/her helpful comments and suggestions.

Data Availability Statement: The manuscript has no associated data.



Arab Journal of Mathematical
Sciences
Emerald Publishing Limited
e-ISSN: 2588-9214
p-ISSN: 1319-5166
DOI 10.1108/AJMS-04-2022-0101

Theorem 1.1. If $f \in L^2(\mathbb{R})$. Then the following are equivalents:

$$(1) \quad \|f(t+h) - f(t)\|_{L^2(\mathbb{R})} = o(h^\delta), \quad (0 < \delta < 1) \text{ as } h \rightarrow 0.$$

$$(2) \quad \int_{|\lambda| \geq r} |\mathcal{F}(f)(\lambda)|^2 d\lambda = o(r^{-2\delta}) \text{ as } r \rightarrow \infty.$$

where $\mathcal{F}(f)$ stands for the Fourier transform of f .

Considerable attention has been devoted to discovering generalizations of new contexts for those theorems, see, e.g. [2–7]. The aim of this paper is to give a generalization of these two theorems by using the harmonic analysis associated with the Laguerre–Bessel operators.

Throughout this paper, C denotes a positive constant which can differ from one line to another.

2. Preliminaries

Given $\alpha \geq 0$. The harmonic analysis on $\mathbb{K} = [0, +\infty[\times [0, +\infty[$ is generated by the following partial differential operators:

$$\begin{cases} \mathcal{D}_{1,\alpha} = \frac{\partial^2}{\partial t^2} + \frac{2\alpha}{t} \frac{\partial}{\partial t} \\ \mathcal{D}_{2,\alpha} = \frac{\partial^2}{\partial x^2} + \frac{2\alpha+1}{x} \frac{\partial}{\partial x} + x^2 \mathcal{D}_{1,\alpha}, \end{cases}$$

where $(x, t) \in \mathbb{K}$. For $(\lambda, m) \in [0, +\infty[\times \mathbb{N}$, the initial value problem:

$$\begin{cases} \mathcal{D}_{1,\alpha} u = -\lambda^2 u, \\ \mathcal{D}_{2,\alpha} u = -4\lambda \left(m + \frac{\alpha+1}{2} \right) u \\ u(0, 0) = 1, \frac{\partial u}{\partial x}(0, 0) = \frac{\partial u}{\partial t}(0, 0) = 0, \end{cases}$$

has a unique solution $\varphi_{\lambda,m}$ given by

$$\varphi_{\lambda,m}(x, t) = j_{\alpha-\frac{1}{2}}(\lambda t) \mathfrak{L}_m^\alpha(\lambda x^2), \quad (x, t) \in \mathbb{K}, \quad (1)$$

where \mathfrak{L}_m^α is the Laguerre function defined on $[0, +\infty[$ by

$$\mathfrak{L}_m^\alpha(x) = e^{-\frac{x}{2}} \frac{L_m^\alpha(x)}{L_m^\alpha(0)}, \quad (2)$$

L_m^α being the Laguerre polynomial of degree m and order α , given by

$$L_m^\alpha(x) = \sum_{k=0}^m (-1)^k \frac{\Gamma(m+\alpha+1)}{\Gamma(k+\alpha+1)} \frac{1}{k!(m-k)!} x^k, \quad (3)$$

and j_α is the normalized Bessel function given by

$$j_\alpha(x) = \Gamma(\alpha+1) \sum_{k=0}^{\infty} \frac{(-1)^k}{k! \Gamma(\alpha+k+1)} \left(\frac{x}{2} \right)^{2k}. \quad (4)$$

Lemma 2.1. [8] For all $(\lambda, m) \in [0, +\infty[\times \mathbb{N}$, the function $\varphi_{\lambda, m}$ is infinitely differentiable on \mathbb{R}^2 , even with respect to each variable and we have

$$\sup_{(x,t) \in \mathbb{K}} |\varphi_{\lambda, m}(x, t)| = 1. \quad (5)$$

Analog of
Titchmarsh's
theorem

Notation. We denote by:

- (1) $|x, t| = |(x, t)|_{\mathbb{K}} = (x^4 + 4t^2)^{\frac{1}{4}}$ the homogeneous norm on \mathbb{K} .
 (2) $|\lambda, m| = |(\lambda, m)|_{[0, +\infty[\times \mathbb{N}} = 4\lambda(m + \frac{\alpha+1}{2})$ the quasinorm on $[0, +\infty[\times \mathbb{N}$. Let us denote \mathbb{B}_r , the ball centered 0 and of radius r , defined by,

$$\mathbb{B}_r = \{(\lambda, m) \in [0, +\infty[\times \mathbb{N}; |\lambda, m| < r\} \quad \text{and} \quad \mathbb{B}_r^c = ([0, +\infty[\times \mathbb{N}) \setminus \mathbb{B}_r.$$

- (3) $L_\alpha^p(\mathbb{K}), p \in [1, +\infty]$, the spaces of measurable functions on \mathbb{K} such that

$$\begin{cases} \|f\|_{p, \alpha} = \left[\int_{\mathbb{K}} |f(x, t)|^p dm_\alpha(x, t) \right]^{\frac{1}{p}} < +\infty, & \text{if } p \in [1, +\infty[\\ \|f\|_{\infty, \alpha} = \text{ess sup}_{(x,t) \in \mathbb{K}} |f(x, t)| < +\infty, \end{cases}$$

where dm_α is the weighted Lebesgue measure on \mathbb{K} , given by

$$dm_\alpha(x, t) = \frac{x^{2\alpha+1} t^{2\alpha}}{\pi \Gamma(\alpha + 1)} dx dt.$$

- (4) $L_{\gamma_\alpha}^p([0, +\infty[\times \mathbb{N}), p \in [1, +\infty]$, the spaces of measurable functions on $[0, +\infty[\times \mathbb{N}$ such that

$$\begin{cases} \|g\|_{\gamma_\alpha, p} = \left[\int_{[0, +\infty[\times \mathbb{N}} |g(\lambda, m)|^p d\gamma_\alpha(\lambda, m) \right]^{\frac{1}{p}} < +\infty, & \text{if } p \in [1, +\infty[\\ \|g\|_{\gamma_\alpha, \infty} = \text{ess sup}_{(\lambda, m) \in [0, +\infty[\times \mathbb{N}} |g(\lambda, m)| < +\infty, \end{cases}$$

where $d\gamma_\alpha$ is the positive measure defined on $[0, +\infty[\times \mathbb{N}$ by

$$\int_{[0, +\infty[\times \mathbb{N}} g(\lambda, m) d\gamma_\alpha(\lambda, m) = \frac{1}{2^{2\alpha-1} \Gamma(\alpha + \frac{1}{2})} \sum_{m=0}^{\infty} L_m^\alpha(0) \int_0^{+\infty} g(\lambda, m) \lambda^{3\alpha+1} d\lambda.$$

Definition 2.2.

- (1) The translation operators $T_{(x,t)}^{(\alpha)}, (x, t) \in \mathbb{K}$ are defined for a continuous function f on \mathbb{K} , by

$$T_{(x,t)}^{(\alpha)} f(y, s) = \begin{cases} \frac{1}{4\pi} \sum_{i,j=0}^1 \int_0^\pi f(\Delta_\theta(x, y), Y + (-1)^i t + (-1)^j s) d\theta, & \text{if } \alpha = 0 \\ b_\alpha \int_{[0, \pi]^3} f(\Delta_\theta(x, y), \Delta_\theta(x, y) \xi) d\mu_\alpha(\xi, \psi, \theta), & \text{if } \alpha > 0. \end{cases}$$

where $\Delta_\theta(x, y) = \sqrt{x^2 + y^2 + 2xy \cos \theta}$, $b_\alpha = \frac{(\alpha+1)\Gamma(\alpha+\frac{1}{2})}{\pi^{\frac{3}{2}}\Gamma(\alpha)}$, $Y = xy \sin \theta$ and

$$d\mu_\alpha(\xi, \psi, \theta) = (\sin \xi)^{2\alpha-1} (\sin \psi)^{2\alpha-1} (\sin \theta)^{2\alpha} d\xi d\psi d\theta.$$

- (2) The convolution product of two continuous functions f, g on \mathbb{K} , with compact support is defined by

$$(f * g)(x, t) = \int_{\mathbb{K}} T_{(x,t)}^{(\alpha)} f(y, s) g(y, s) dm_\alpha(y, s), \quad (x, t) \in \mathbb{K}.$$

We have the following properties:

- (1) If $f \in L_\alpha^p(\mathbb{K}), g \in L_\alpha^q(\mathbb{K})$ such that $1 \leq p, q \leq \infty$ and $\frac{1}{p} + \frac{1}{q} - 1 = \frac{1}{r}$ then the function $f * g \in L_\alpha^r(\mathbb{K})$, and

$$\|f * g\|_{r,\alpha} \leq \|f\|_{p,\alpha} \|g\|_{q,\alpha}.$$

- (2) For all $(\lambda, m) \in [0, +\infty[\times \mathbb{N}$, the kernel $\varphi_{\lambda,m}$ verifies the following product formula

$$\varphi_{\lambda,m}(x, t) \varphi_{\lambda,m}(y, s) = T_{(x,t)}^{(\alpha)} \varphi_{\lambda,m}(y, s), \quad (x, t), (y, s) \in \mathbb{K}.$$

- (3) For $f \in L_\alpha^p(\mathbb{K}), p \in [1, +\infty]$, we have $T_{(x,t)}^{(\alpha)} f \in L_\alpha^p(\mathbb{K})$ and

$$\|T_{(x,t)}^{(\alpha)} f\|_{p,\alpha} \leq \|f\|_{p,\alpha}.$$

The Fourier–Laguerre–Bessel transform of a function in $L_\alpha^1(\mathbb{K})$ is given by

$$\mathcal{F}_{LB} f(\lambda, m) = \int_{\mathbb{K}} f(x, t) \varphi_{\lambda,m}(x, t) dm_\alpha(x, t), \quad (\lambda, m) \in [0, +\infty[\times \mathbb{N}.$$

From Ref. [8], it is well known that Fourier–Laguerre–Bessel transform can be inverted to

$$\mathcal{F}_{LB}^{-1} f(x, t) = \int_{[0, +\infty[\times \mathbb{N}} f(\lambda, m) \varphi_{\lambda,m}(x, t) d\gamma_\alpha(\lambda, m), \quad (x, t) \in \mathbb{K}.$$

It is well-known (see Refs. [8–11]) that the Fourier–Laguerre–Bessel transform \mathcal{F}_{LB} satisfies the following properties.

Theorem 2.3. (Inversion formula). If $f \in L_\alpha^1(\mathbb{K})$ such that $\mathcal{F}_{LB}(f) \in L_{\gamma_\alpha}^1([0, +\infty[\times \mathbb{N})$, then for all $(x, t) \in \mathbb{K}$ we have

$$f(x, t) = \int_{[0, +\infty[\times \mathbb{N}} \mathcal{F}_{LB} f(\lambda, m) \varphi_{\lambda,m}(x, t) d\gamma_\alpha(\lambda, m).$$

Theorem 2.4. (Plancherel Theorem for \mathcal{F}_{LB}). The generalized Fourier transform \mathcal{F}_{LB} extends to an isometric isomorphism from $L_\alpha^2(\mathbb{K})$. Onto $L_{\gamma_\alpha}^2([0, +\infty[\times \mathbb{N})$.

Proposition 2.5. For $f \in L_\alpha^1(\mathbb{K}), (x, t) \in \mathbb{K}$ and $(\lambda, m) \in [0, +\infty[\times \mathbb{N}$, we have

$$\mathcal{F}_{LB} \left(T_{(x,t)}^{(\alpha)} f \right) (\lambda, m) = \varphi_{\lambda,m}(x, t) \mathcal{F}_{LB}(f)(\lambda, m). \quad (6)$$

Remark 1. From (6) (see Ref. [14]), we get

$$\mathcal{F}_{LB}\left(T_{(x,t)}^{(\alpha)}f - f\right)(\lambda, m) = (\varphi_{\lambda,m}(x, t) - 1)\mathcal{F}_{LB}(f)(\lambda, m). \quad (7)$$

Analog of
Titchmarsh's
theorem

3. Main results

In order to give the main results, we begin with auxiliary results interesting in themselves.

Lemma 3.1. Let $\eta > 0$.

(1) The behavior in 0 of the kernel $\varphi_{\lambda,m}$ could be expressed as follows:

$$\varphi_{\lambda,m}(x, t) = 1 - \frac{(\lambda t)^2}{4(\alpha + \frac{1}{2})} - \frac{|\lambda, m| x^2}{4(\alpha + 1)} + \kappa_{\alpha,m} \lambda^2 x^4 + o\left(|\lambda, m|^2 |x, t|^4\right), \quad (8)$$

where $\kappa_{\alpha,m} = \frac{m^2}{2(\alpha+1)(\alpha+2)} + \frac{m}{2(\alpha+2)} + \frac{1}{8}$

(2) There exists a constant C , such that if $|\lambda, m| x^2 < \eta$, then

$$|\varphi_{\lambda,m}(x, t) - 1| \geq C|\lambda, m| x^2. \quad (9)$$

(3) There exist $C > 0$ such that for all $(x, t) \in \mathbb{K}$,

$$|\lambda, m| |x, t|^2 < \eta \Rightarrow |\varphi_{\lambda,m}(x, t) - 1|^2 \leq C|\lambda, m|^2 |x, t|^4 \quad (10)$$

(4) There exist $C > 0$ and $A > 0$ such that for all $|x, t|^2 |\lambda, m| > A$ and $(x, t) \in \mathbb{K}$,

$$|\varphi_{\lambda,m}(x, t) - 1| \geq C. \quad (11)$$

Proof.

(1) From the relation (2) and (3), we have

$$\mathfrak{L}_m^\alpha(x) = 1 - \frac{m + \frac{\alpha+1}{2}}{\alpha + 1} x + \left(\frac{m^2}{2(\alpha + 1)(\alpha + 2)} + \frac{m}{2(\alpha + 2)} + \frac{1}{8} \right) x^2 + o(x^2). \quad (12)$$

Then (i) could be deduced easily using the relation (1),(12) and the behavior in 0 of the normalized Bessel function which states

$$j_\alpha(u) = 1 - \frac{1}{4(\alpha + 1)} u^2 + o(u^2).$$

(2) Using relations (8), we obtain

$$\lim_{|\lambda, m| x^2 \rightarrow 0} \left(\frac{|\varphi_{\lambda,m}(x, t) - 1|}{|\lambda, m| x^2} \right) = \frac{1}{4(\alpha + 1)} > 0,$$

which proves the wanted result.

(3) Using relation (8).

(4) From ([6], Lemma 4.3), we have

$$\lim_{|\lambda, m| \rightarrow +\infty} \psi_{\lambda, m}(x, t) = 0,$$

where $\psi_{\lambda, m}(x, t) = e^{i\lambda t} \mathfrak{L}_m^\alpha(\lambda x^2)$ the Laguerre kernel, and from Ref. [12], we have the asymptotic formula for the normalized Bessel function j_α when $x \rightarrow +\infty$:

$$j_\alpha(x) = \frac{\Gamma(\alpha+1)}{\Gamma(\frac{1}{2})} \left(\frac{2}{x}\right)^{\alpha+\frac{1}{2}} \cos\left(x - (2\alpha+1)\frac{\pi}{4}\right) + o\left(\frac{1}{x^{\frac{3}{2}}}\right).$$

Hence as

$$\varphi_{\lambda, m}(x, t) = j_{\alpha-\frac{1}{2}}(\lambda t) \frac{1}{e^{i\lambda t}} \psi_{\lambda, m}(x, t),$$

then $\lim_{|\lambda, m| \rightarrow +\infty} \varphi_{\lambda, m}(x, t) = 0$, we get $\lim_{|\lambda, m| \rightarrow +\infty} |\varphi_{\lambda, m}(x, t) - 1| = 1$, which completes the proof. \square

Lemma 3.2. (Hausdorff-Young inequality) Let $1 < p \leq 2$. If $f \in L_\alpha^p(\mathbb{K})$, then $\mathcal{F}_{LB}f \in L_{\gamma_\alpha}^q([0, +\infty[\times \mathbb{N})$ and we have

$$\|\mathcal{F}_{LB}f\|_{\gamma_{\alpha, q}} \leq C \|f\|_{p, \alpha},$$

where the numbers p and q above are conjugate exponents:

$$\frac{1}{p} + \frac{1}{q} = 1.$$

Proof. By applying the Riesz–Thorin interpolation theorem to the elementary estimate [13] and Plancherel theorem, we obtain the desired inequality. \square

Proposition 3.3. Let f be a function in $L_\alpha^p(\mathbb{K})$, such that $\left\| T_{(x, t)}^{(\alpha)} f - f \right\|_{p, \alpha} = O(x^\gamma)$ for $1 < p \leq 2$ and $0 < \gamma \leq 1$. Then $\mathcal{F}_{LB}f$ belongs to $L_{\gamma_\alpha}^p([0, +\infty[\times \mathbb{N})$, where

$$\frac{(\alpha+2)p}{(\alpha+2)(p-1) + \frac{p}{2}} < \beta \leq \frac{p}{p-1}.$$

Proof. By proceeding similarly to theorem (Theorem 3.1 [6]). For fixed $(x, t) \in \mathbb{K}$, we have using relations (7) and Lemma 3.2

$$\int_{[0, +\infty[\times \mathbb{N}} |\varphi_{\lambda, m}(x, t) - 1|^q |\mathcal{F}_{LB}f(\lambda, m)|^q d\gamma_\alpha(\lambda, m) = O(x^{\gamma q}).$$

Using relations (9), we get

$$\begin{aligned} \int_{\mathbb{B}_{\frac{\eta}{x^2}}} |\lambda, m|^q |\mathcal{F}_{LB}f(\lambda, m)|^q d\gamma_\alpha &\leq x^{-2q} \int_{\mathbb{B}_{\frac{\eta}{x^2}}} |\varphi_{\lambda, m}(x, t) - 1|^q |\mathcal{F}_{LB}f(\lambda, m)|^q d\gamma_\alpha \\ &\leq C x^{(\gamma-2)q}. \end{aligned}$$

Now, let $\beta \leq q$. From Hölder inequality, one gets

$$\int_{\mathbb{B}_X} |\lambda, m|^\beta |\mathcal{F}_{LB}f(\lambda, m)|^\beta d\gamma_\alpha = \left(\int_{\mathbb{B}_X} |\lambda, m|^q |\mathcal{F}_{LB}f(\lambda, m)|^q d\gamma_\alpha \right)^{\frac{\beta}{q}} \left(\int_{\mathbb{B}_X} 1 d\gamma_\alpha \right)^{1-\frac{\beta}{q}}.$$

Analogue of
Titchmarsh's
theorem

Therefore

$$\int_{\mathbb{B}_X} |\lambda, m|^\beta |\mathcal{F}_{LB}f(\lambda, m)|^\beta d\gamma_\alpha(\lambda, m) = O\left(X^{\frac{(2-\gamma)q}{2}\frac{\beta}{q} + (\alpha+2)(1-\frac{\beta}{q})}\right). \quad (13)$$

Recall that $\mathbb{B}_1^c = ([0, +\infty[\times \mathbb{N}) \setminus \mathbb{B}_1$. To get the theorem, it is enough to prove that $\int_{\mathbb{B}_1^c \cap \mathbb{B}_X} |\mathcal{F}_{LB}f(\lambda, m)|^\beta d\gamma_\alpha(\lambda, m)$ is bounded when $X \rightarrow +\infty$. Therefore, we can write

$$\int_{\mathbb{B}_1^c \cap \mathbb{B}_X} |\mathcal{F}_{LB}f(\lambda, m)|^\beta d\gamma_\alpha(\lambda, m) = \frac{1}{2^{2\alpha-1}\Gamma(\alpha + \frac{1}{2})} \sum_{m=0}^{+\infty} L_m^\alpha(0)I,$$

where I depend on m and X and has the expression

$$I = \int_{\frac{1}{4m+2\alpha+2}}^{\frac{X}{4m+2\alpha+2}} |\mathcal{F}_{LB}f(\lambda, m)|^\beta \lambda^{3\alpha+1} d\lambda.$$

Then

$$I = \int_{\frac{1}{4m+2\alpha+2}}^{\frac{X}{4m+2\alpha+2}} (4m+2\alpha+2) |\lambda, m|^{-\beta} \Phi_m'(|\lambda, m|) d\lambda,$$

where

$$\Phi_m(X) = \int_{\frac{1}{4m+2\alpha+2}}^{\frac{X}{4m+2\alpha+2}} |\lambda, m|^\beta |\mathcal{F}_{LB}f(\lambda, m)|^\beta \lambda^{3\alpha+1} d\lambda$$

Making a change of variables and an integration by parts, we get

$$I = \Phi_m(X)X^{-\beta} + \beta \int_1^X t^{-\beta-1} \Phi_m(t) dt.$$

Consequently

$$\int_{\mathbb{B}_1^c \cap \mathbb{B}_X} |\mathcal{F}_{LB}f(\lambda, m)|^\beta d\gamma_\alpha(\lambda, m) = X^{-\beta} \psi(X) + \beta \int_1^X t^{-\beta-1} \psi(t) dt,$$

where

$$\begin{aligned} \psi(X) &= \frac{1}{2^{2\alpha-1}\Gamma\left(\alpha + \frac{1}{2}\right)} \sum_{m=0}^{+\infty} L_m^\alpha(0) \Phi_m(X), \\ &= \int_{\mathbb{B}_1^c \cap \mathbb{B}_X} |\lambda, m|^\beta |\mathcal{F}_{LB}f(\lambda, m)|^\beta d\gamma_\alpha(\lambda, m). \end{aligned}$$

From relation (13), we have

$$\int_{\mathbb{B}_1^c \cap \mathbb{B}_X} |\mathcal{F}_{LB}f(\lambda, m)|^\beta d\gamma_\alpha(\lambda, m) = O\left(X^{-\beta + \frac{2-\gamma}{2}\beta + (\alpha+2)(1-\frac{\beta}{q})}\right) \\ + O\left(\int_1^X t^{-\beta-1} t^{\frac{2-\gamma}{2}\beta + (\alpha+2)(1-\frac{\beta}{q})} dt\right).$$

This is bounded as $X \rightarrow +\infty$ if $-\beta\left(\frac{\gamma}{2} + \frac{\alpha+2}{q}\right) + (\alpha+2) < 0$ that gives $\beta > \frac{(\alpha+2)p}{(\alpha+2)(p-1) + \frac{p}{\gamma}}$. \square

Next we define the ψ -Laguerre–Bessel–Lipschitz class:

Definition 3.4. A function f is said to be in ψ -Laguerre–Bessel–Lipschitz class and is denoted by $Lip_\alpha(\psi, 2)$, if f belongs to $L_\alpha^2(\mathbb{K})$ and verifies, for all $(x, t) \in \mathbb{K}$

$$\left\| T_{(x,t)}^{(\alpha)} f - f \right\|_{2,\alpha} = O(\psi(|x, t|)) \text{ as } |x, t| \rightarrow 0, \quad (14)$$

where

- (1) $\psi(t)$ is a continuous increasing function on $[0; \infty[$.
- (2) $\psi(0) = 0$ and $\psi(ts) = \psi(t)\psi(s)$ for all $t, s \in [0; \infty[$.
- (3) $\int_0^h s\psi(s^{-1})ds = o\left(\frac{1}{h^2}\psi(h)\right)$ as $h \rightarrow 0$.

Example 3.5. Let $\psi(t) = t^\gamma$, where $0 < \gamma < 1$. In this case, the relation (14) is a generalization of Lipschitz condition $\|f(x+h) - f(x)\| = O(h^\gamma)$, and the ψ -Laguerre–Bessel–Lipschitz class $Lip_\alpha(\psi, 2)$, are called the Laguerre–Bessel–Lipschitz class $Lip_\alpha(\gamma, 2)$.

Now, we are able to generalise the equivalence theorem.

Theorem 3.6. Let $f \in L_\alpha^2(\mathbb{K})$, the following two conditions are equivalent:

- (1) $f \in Lip_\alpha(\psi, 2)$.
- (2)

$$\int_{\mathbb{B}_r^c} |\mathcal{F}_{LB}f(\lambda, m)|^2 d\gamma_\alpha(\lambda, m) = o(\psi(r^{-1})) \text{ as } r \rightarrow +\infty.$$

Proof. (i \Rightarrow ii): Let $f \in L_\alpha^2(\mathbb{K})$ from (i), we have

$$\left\| T_{(x,t)}^{(\alpha)} f - f \right\|_{2,\alpha}^2 = \int_{[0,+\infty[\times \mathbb{N}} |\varphi_{\lambda,m}(x, t) - 1|^2 |\mathcal{F}_{LB}f(\lambda, m)|^2 d\gamma_\alpha.$$

Therefore, using relation (11), we have

$$\int_{\frac{\mathbb{A}}{|x,t|^2}} |\mathcal{F}_{LB}f(\lambda, m)|^2 d\gamma_\alpha \leq C \int_{\frac{\mathbb{A}}{|x,t|^2}} |\varphi_{\lambda,m}(x, t) - 1|^2 |\mathcal{F}_{LB}f(\lambda, m)|^2 d\gamma_\alpha \\ \leq C \left\| T_{(x,t)}^{(\alpha)} f - f \right\|_{2,\alpha}^2 = O(\psi(|x, t|^2)).$$

Consequently, (ii) holds.

(ii \Rightarrow i): Denote $r = \frac{\eta}{|x,t|^2}$, by Plancherel theorem, we get

$$\left\| T_{(x,t)}^{(\alpha)} f - f \right\|_{2,\alpha}^2 = I_1 + I_2,$$

where

$$I_1 = \int_{\mathbb{B}_r} |\varphi_{\lambda,m}(x,t) - 1|^2 |\mathcal{F}_{LB} f(\lambda,m)|^2 d\gamma_\alpha(\lambda,m)$$

and

$$I_2 = \int_{\mathbb{B}_r^c} |\varphi_{\lambda,m}(x,t) - 1|^2 |\mathcal{F}_{LB} f(\lambda,m)|^2 d\gamma_\alpha(\lambda,m).$$

Using relation (5), we find that

$$I_2 \leq 4 \int_{\mathbb{B}_r^c} |\mathcal{F}_{LB} f(\lambda,m)|^2 d\gamma_\alpha = o(\psi(r^{-1})) = o(\psi(|x,t|^2)).$$

Denote

$$g(X) = \int_X^\infty |\mathcal{F}_{LB} f(\lambda,m)|^2 \lambda^{3\alpha+1} d\lambda,$$

then $g'(\lambda) = -|\mathcal{F}_{LB} f(\lambda,m)|^2 \lambda^{3\alpha+1}$. Using relation (10), which gives

$$I_1 \leq C|x,t|^4 \frac{1}{2^{2\alpha-1}\Gamma(\alpha+\frac{1}{2})} \sum_{m=0}^\infty L_m^\alpha(0) \left(\int_0^{\frac{\eta}{4\kappa_m|x,t|^2}} (4\kappa_m)^2 \lambda^2 (-g'(\lambda)) d\lambda \right),$$

by integration by parts, we have

$$I_1 \leq C|x,t|^4 \frac{1}{2^{2\alpha-1}\Gamma(\alpha+\frac{1}{2})} \sum_{m=0}^\infty L_m^\alpha(0) \left(-\frac{\eta^2}{|x,t|^4} g\left(\frac{\eta}{4\kappa_m|x,t|^2}\right) + (4\kappa_m)^2 \int_0^{\frac{\eta}{4\kappa_m|x,t|^2}} 2\lambda g(\lambda) d\lambda \right).$$

Remark that

$$\frac{1}{2^{2\alpha-1}\Gamma(\alpha+\frac{1}{2})} \sum_{m=0}^\infty L_m^\alpha(0) g\left(\frac{R}{4\kappa_m}\right) = \int_{\mathbb{B}_R^c} |\mathcal{F}_{LB} f(\lambda,m)|^2 d\gamma_\alpha = o(\psi(R^{-1})).$$

Making a change of variable, one gets

$$\begin{aligned} I_1 &\leq o(\psi(|x,t|^2)) + C|x,t|^4 \int_0^{\frac{\eta}{|x,t|^2}} u \frac{1}{2^{2\alpha-1}\Gamma(\alpha+\frac{1}{2})} \sum_{m=0}^\infty L_m^\alpha(0) g\left(\frac{u}{4\kappa_m}\right) du \\ &= o(\psi(|x,t|^2)) + |x,t|^4 o\left(\int_0^{\frac{\eta}{|x,t|^2}} u \psi(u^{-1}) du\right) \\ &= o(\psi(|x,t|^2)). \end{aligned}$$

$$\left\| T_{(x,t)}^{(a)} f - f \right\|_{2,a}^2 = o\left(\psi(|x,t|^2)\right) = o\left(\psi(|x,t|)^2\right) \quad \text{as } |x,t| \rightarrow 0.$$

□

We conclude this work by the following immediate consequence. It is analogous of Titchmarsh theorem ([1], Theorem 85) is established for Laguerre–Bessel transform.

Corollary 3.7. *Let $\psi(t) = t^\gamma$, where $0 < \gamma < 1$. The following two conditions are equivalent:*

- (1) *f is in Laguerre–Bessel–Lipschitz class $Lip_\alpha(\gamma, 2)$.*
- (2) $\int_{\mathbb{R}_+^c} |\mathcal{F}_{LB} f(\lambda, m)|^2 d\gamma_\alpha(\lambda, m) = O(r^{-\gamma})$ as $r \rightarrow +\infty$.

References

1. Titchmarsh EC. Introduction to the theory of fourier integral. Amen House, London. E. C. 4: Oxford University Press; 1948.
2. El Ouadih S, Daher R. Characterization of Dini-Lipschitz functions for the Helgason Fourier transform on rank one symmetric spaces. Adv Pure Appl Math. 2016; 7(4): 223-30.
3. Daher R, El Hamma M. An analog of Titchmarsh's theorem for the generalized Dunkl transform. J.Pseud Diff Op Appl. 2016; 7(1): 59-65.
4. Younis MS. Fourier transform of Lipschitz functions on compact groups. Ph.D. Thesis. Hamilton, ON, Canada: McMaster University; 1974.
5. El ouadih S, Daher R. Generalization of Titchmarsh's theorem for the Dunkl transform in the space $Lp(\mathbb{R}^d; w_l(x)dx)$.
6. Negzaoui S. Lipschitz conditions in Laguerre hypergroup. Mediterr J Math. 2017; 14: 191. doi: [10.1007/s00009-017-0989-4](https://doi.org/10.1007/s00009-017-0989-4).
7. Achak A, Bouhlal A, Daher R. Titchmarsh's theorem and some remarks concerning the right-sided quaternion Fourier transform. Bol Soc Mat Mex. 2020; 26: 599-616.
8. Jebbari E, Sifi M, Soltani F. Laguerre-Bessel wavelet transform. Glob J Pure Appl. Math. 2005; 1: 13-26.
9. Kortas H, Sifi M. Levy-Khintchine formula and dual convolution semigroups associated with Laguerre and Bessel functions. Potential Anal. 2001; 15: 43-58.
10. Jebbari E. Harmonic analysis associated with system of partial differential operators D_1 and D_2 . Preprint, 2004.
11. Soumeia H, Lotfi K. Uncertainty principle inequalities related to Laguerre-Bessel transform.
12. Watson GN. A treatise on the theory of Bessel functions. Cambridge: Cambridge University Press; 1966.
13. Folland G. Real analysis modern techniques and their applications, Pure Appl. Math. 2nd ed. New York: Wiley; 1999.
14. Rakhimi Larbi, Daher Radouan. Modulus of Smoothness and K-Functionals Constructed by Generalized Laguerre-Bessel Operator. Tatra Mountains Mathematical Publications. 2022; 81(1). doi: [10.2478/tmmp-2022-0008](https://doi.org/10.2478/tmmp-2022-0008).

Corresponding author

Larbi Rakhimi can be contacted at: larbi.rakhimi-etu@etu.univh2c.ma

For instructions on how to order reprints of this article, please visit our website:

www.emeraldgroupublishing.com/licensing/reprints.htm

Or contact us for further details: permissions@emeraldinsight.com